

# Treatment with Rutin - A Therapeutic Strategy for Neutrophil-Mediated Inflammatory and Autoimmune Diseases

## - Anti-inflammatory Effects of Rutin on Neutrophils -

Bahareh Abd Nikfarjam<sup>1,2</sup>, Mohtaram Adineh<sup>2</sup>, Farid Hajiali<sup>1</sup>, Marjan Nassiri-Asl<sup>1,3\*</sup>

<sup>1</sup>Cellular and Molecular Research Center, Qazvin University of Medical Sciences, Qazvin, Iran

<sup>2</sup>Department of Immunology, Qazvin University of Medical Sciences, Qazvin, Iran

<sup>3</sup>Department of Pharmacology, School of Medicine, Qazvin University of Medical Sciences, Qazvin, Iran

### Key Words

human neutrophils, myeloperoxidase, nitric oxide, rutin, tumor necrosis factor- $\alpha$

### Abstract

**Objectives:** Neutrophils represent the front line of human defense against infections. Immediately after stimulation, neutrophilic enzymes are activated and produce toxic mediators such as pro-inflammatory cytokines, nitric oxide (NO) and myeloperoxidase (MPO). These mediators can be toxic not only to infectious agents but also to host tissues. Because flavonoids exhibit antioxidant and anti-inflammatory effects, they are subjects of interest for pharmacological modulation of inflammation. In the present study, the effects of rutin on stimulus-induced NO and tumor necrosis factor (TNF)- $\alpha$  productions and MPO activity in human neutrophils were investigated.

**Methods:** Human peripheral blood neutrophils were isolated using Ficoll-Hypaque density gradient centrifugation coupled with dextran T500 sedimentation. The cell preparations containing > 98% granulocytes were determined

by morphological examination through Giemsa staining. Neutrophils were cultured in complete Roswell Park Memorial Institute (RPMI) medium, pre-incubated with or without rutin (25  $\mu$ M) for 45 minutes, and stimulated with phorbol 12-myristate 13-acetate (PMA). Then, the TNF- $\alpha$ , NO and MPO productions were analyzed using enzyme-linked immunosorbent assay (ELISA), Griess Reagent, and MPO assay kits, respectively. Also, the viability of human neutrophils was assessed using tetrazolium salt 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyl tetrazolium bromide (MTT), and neutrophils were treated with various concentrations of rutin (1 - 100  $\mu$ M), after which MTT was appended and incubated at 37°C for 4 hour.

**Results:** Rutin at concentrations up to 100  $\mu$ M did not affect neutrophil viability during the 4-hour incubation period. Rutin significantly decreased the NO and TNF- $\alpha$  productions in human peripheral blood neutrophils compared to PMA-control cells ( $P < 0.001$ ). Also, MPO activity was significantly reduced by rutin ( $P < 0.001$ ).

**Conclusion:** In this *in vitro* study, rutin had an anti-inflammatory effect due to its inhibiting NO and TNF- $\alpha$  productions, as well as MPO activity, in activated human neutrophils. Treatment with rutin may be considered as a therapeutic strategy for neutrophil-mediated inflammatory/ autoimmune diseases.

Received: Jan 04, 2017 Reviewed: Feb 01, 2017 Accepted: Feb 03, 2017

© This is an Open-Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted noncommercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

© This paper meets the requirements of KS X ISO 9706, ISO 9706-1994 and ANSI/NISO Z39.48-1992 (Permanence of Paper).

\*Corresponding Author

Marjan Nassiri-Asl. Cellular and Molecular Research Centre, Department of Pharmacology, Qazvin University of Medical Sciences, Qazvin, PO 341197-5981, Iran.

Tel: +98-283-333-6001 Fax: +98-283-332-4970

E-mail: [marjannassiriaslm@gmail.com](mailto:marjannassiriaslm@gmail.com)

© 2017 Korean Pharmacopuncture Institute

<http://www.journal.ac>